

JKI Data Sheets

Plant Diseases and Diagnosis

Venche TALGØ / Gary CHASTAGNER

Phytophthora on *Abies* spp. (true firs)



Imprint

The open access series “JKI Data Sheets – Plant Diseases and Diagnosis” is a publication that publishes original papers, pathogen descriptions, findings and reports on biotic and abiotic causes of crop diseases and crop damage.

All manuscripts submitted for publication in the JKI Data Sheets are peer-reviewed by at least two independent referees while the anonymity of author(s) is preserved.

All contributions are made available under the Creative Commons licence. This allows you to use and distribute the whole work or parts of the work at no charge as long as you use it only for noncommercial purposes, name the author(s) and source(s) and do not modify the work.

Publisher/Editor-in-Chief: Dr. Georg F. Backhaus, Präsident und Professor
Julius Kühn-Institut, Federal Research Centre for Cultivated Plants
Erwin-Baur-Str. 27
D-06484 Quedlinburg
Germany

Managing Editor: Dr. Olaf Hering, Information Centre and Library
Julius Kühn-Institut
Königin-Luise-Str. 19
D-14195 Berlin
Germany
redaktion.datasheets@jki.bund.de

Submission of manuscripts: Please go to the journal's website at
<http://pub.jki.bund.de/>

ISSN: 2191-1398

DOI: 10.5073/jkidsppd.2013.077

Importance of *Abies* spp.

Approximately 50 species belong to the genus *Abies* and they are mainly native to the temperate regions of the northern hemisphere (<http://www.discoverlife.org>). They play a very important environmental role in their regions of origin. Several species are also important for timber production outside their native range.

Others are highly appreciated as ornamentals in landscape plantings. Because of their natural conical shape, color and strong branches for holding ornaments, true firs, especially species with excellent postharvest needle retention, are also ideally suited for use as Christmas trees and bough production. In Europe, the most common species grown as Christmas trees is Nordmann fir, while in the USA, Fraser (*Abies fraseri*) and noble fir (*Abies procera*) predominate.

Other *Abies*-species marketed as Christmas trees include balsam fir (*A. balsamea*) grand fir (*A. grandis*), Korean fir (*Abies koreana*), Siberian fir (*A. sibirica*), subalpine fir (*A. lasiocarpa*), Turkish fir (*A. bornmuelleriana*), and white fir (*A. concolor*). Noble fir is the main species for bough production both in Europe and USA.

Phytophthora species

A number of *Phytophthora* spp. has been isolated from true firs worldwide. The following is a partial list of species that have been reported on specific hosts:

<i>Phytophthora</i> species	Disease symptoms	<i>Abies</i> species	References
<i>cactorum</i>	root rot stem canker branch flagging	<i>amabilis</i> <i>balsamea</i> <i>balsamea</i> var. <i>phanerolepis</i> <i>concolor</i> <i>fraseri</i> <i>magnifica</i> var. <i>shastensis</i> <i>procera</i>	Adams & Bielenin, 1988 Chastagner <i>et al.</i> , 1995 Hamm & Hansen, 1982 Hong & Marston, 2005
<i>cambivora</i>	root rot stem canker branch flagging	<i>procera</i>	Chastagner <i>et al.</i> , 1995 Talgø <i>et al.</i> , 2006
<i>capsici</i>		<i>fraseri</i>	Quesada-Ocampo <i>et al.</i> , 2009
<i>cinnamomi</i>	root rot stem canker branch flagging	<i>fraseri</i> <i>procera</i>	Chastagner <i>et al.</i> , 1995 Grand & Lapp, 1974
<i>citricola</i>	root rot shoot blight	<i>balsamea</i> <i>concolor</i> <i>fraseri</i> <i>magnifica</i> <i>procera</i>	Adams & Bielenin, 1988 McCain & Scharpf, 1986 Shew & Benson, 1981
<i>citrophthora</i>	root rot collar rot	<i>lasiocarpa</i> var. <i>arizonica</i>	Józsa <i>et al.</i> , 2011
<i>cryptogea</i>	root rot stem canker branch flagging	<i>procera</i>	Chastagner <i>et al.</i> , 1995
<i>drechsleri</i>	root rot stem canker branch flagging	<i>fraseri</i> <i>procera</i>	Benson <i>et al.</i> , 1976 Chastagner <i>et al.</i> , 1995
<i>gonapodyides</i>	root rot stem canker branch flagging	<i>procera</i>	Chastagner <i>et al.</i> , 1995
<i>inundata</i> -like		<i>nordmanniana</i>	Talgø <i>et al.</i> , 2007
<i>megasperma</i>	root rot stem canker	<i>lasiocarpa</i> <i>procera</i>	Chastagner <i>et al.</i> , 1995 Talgø <i>et al.</i> , 2007
<i>pseudotsugae</i>	root rot stem canker branch flagging	<i>procera</i>	Chastagner <i>et al.</i> , 1995
<i>ramorum</i> ¹	root rot stem canker	<i>concolor</i> <i>magnifica</i> <i>grandis</i>	Chastagner & Riley, 2010 Riley <i>et al.</i> , 2011

¹ in the European Union *P. ramorum* is a regulated organism (see chapter 'Quarantine recommendation')

Disease symptoms (see figures)

Wet conditions are necessary for the spread of *Phytophthora* spores and infection of plants. Most *Phytophthora* spp. are soil borne, which means that spores are spread in water films in infested soil and may cause root rot and possibly spread to stems in the form of cankers. The first visible symptoms caused by soil borne *Phytophthora* spp. can include a general discoloration of foliage (chlorotic, greyish, and finally necrotic) or one or more dying basal branches (flagging). The foliage discoloration is due to lack of water and nutrients as the pathogen destroys the vascular system.

Flagging indicates that an aboveground canker has developed, commonly extending upwards on one side of the tree and killing the branches in its path. Multiple cankers and/or spiral development of a single canker may girdle the tree. The cankers appear slightly sunken, sometimes cracked along the margins, and darker than the healthy bark. Heavy resin flow may occur. Beneath the bark of the cankered areas, the tissue appears brick red with a sharp border at the healthy, green tissue along the leading edges. Symptoms caused by *P. cambivora* on noble fir in Norway are presented in Figure 1.

In some cases, especially in nurseries, lower foliage may be infected when water splashes contaminated soil onto the foliage. The use of contaminated irrigation water may cause shoot blight, which is also a common symptom caused by airborne *Phytophthora* spp. like *P. ramorum*. The presence of *P. ramorum* spores during bud break and shoot elongation may also cause infection and subsequent canker development on branches of susceptible fir species.

Possibility of symptom confusion

Both Armillaria root rot (*Armillaria* spp.) and Annosus root rot (*Heterobasidion annosum*) may kill the roots of true firs, resulting in discolored foliage similar to the symptoms caused by *Phytophthora* spp., but the two pathogens produce clear signs that distinguish the diseases they cause from *Phytophthora* root rot. *Armillaria* spp. produce root-like rhizomorphs that are visible on the roots and root collar, white mycelial fans beneath the bark near the root collar, and fruiting bodies at stem bases.

H. annosum produces a thin, colorful (pink, purple, red, brown) mycelial layer between the bark and the wood as well as conks on root collars. The conks are often hidden by litter, but a typical, irregularly-shaped dark stain in the older wood of the trunk is normally a reliable symptom and sufficient for diagnosing the pathogen.

Disease development

Damage potential by *Phytophthora* spp. on Christmas trees is considered moderate to high (Chastagner & Byther, 1997). *Phytophthora* spp. produce motile spores under wet conditions. They can actively swim for a few millimeters in water, enough to reach and infect nearby roots, or be transported over longer distances in windborne droplets or by water movement in soil or along waterways.

Soil borne *Phytophthora* spp. can spread up the roots and kill the cambium and inner bark. Disease development depends on several factors; health of the seedling transplants, soil type, weather conditions, and the *Phytophthora* and *Abies* species in question. Disease-free nursery stock is essential for a good outcome. Plants grown in well-drained soils, as opposed to poorly-drained heavy soils, are less prone to *Phytophthora* problems. In cases where transplants are latently infected (e.g. where the pathogen is present, but symptoms are suppressed by fungicides), and then are planted in saturated soil, mortality can be very high in a short period of time (Talgø *et al.*, 2007).

Another concern is that infected transplants may introduce the pathogen into uninfested fields. The aggressiveness of different *Phytophthora* spp. varies, and some *Abies* spp. are more susceptible than others. In North Carolina (USA), *P. cinnamomi* is a well-known pathogen on native Fraser fir that is grown in Christmas tree plantations (Grand & Lapp, 1974). In the western USA, noble fir grows naturally in the mountain areas and has also become the major Christmas tree grown in plantations in the region. In Oregon and Washington State a total of eight *Phytophthora* spp. are associated with root rot development on noble fir, with *P. cactorum*, *P. cambivora*, *P. cinnamomi*, and *P. cryptogea* considered to be the most aggressive species.

Laboratory and field studies in North Carolina and in the U.S. Pacific Northwest have clearly demonstrated that there are significant differences in the susceptibility of true firs to *Phytophthora* root rot (Bensen *et al.*, 1997; Chastagner, 2010; Cooley *et al.*, 1988; Frampton & Benson, 2012; Hinesley *et al.*, 2000) (Fig. 2). Variation in susceptibility is dependent on the *Phytophthora* spp, host, and environmental conditions. Noble, balsam (*A. balsamea*), grand, California red (*Abies magnifica*), Fraser, and Shasta (*A. magnifica* var. *shastensis*) firs are among the more susceptible species, whereas Turkish, European silver (*A. alba*), Veitch (*A. veitchii*), Nordmann, Momi (*A. firma*), and Korean firs tend to be much less susceptible.

Diagnosis

It is not possible to identify a *Phytophthora* infection only by disease symptoms. Different diagnostic techniques like direct isolation, molecular and serological methods help to identify *Phytophthora* as the cause of the tree disease and to specify the *Phytophthora* species. Information on *Phytophthora* diagnosis on trees is given for example in <http://forestphytophthoras.org/key-to-species>, <http://www.phytophthoradb.org>, <http://phytophthora-id.org> and in Martin *et al.* (2012).

Please contact your national authorities (see next chapter) for help with diagnosis.

What to do in case trees are suspected to be infected?

Contact your responsible national authorities, for example: [addresses.pdf](#)

Management and control

The use of healthy transplant seedlings and site selection are the most important cultural factors affecting the management of *Phytophthora* root rot. At the farm level, proper drainage is required to avoid saturated soils. Highly susceptible firs should not be planted on sites with heavy, poorly drained soils. Avoiding the use of overhead irrigation, particularly if the water is contaminated with *Phytophthora* inoculum, will prevent shoot blight.

Pruning should not be done during wet periods to enable a quick drying of the wound and a quick periderm development. Injury of the stem, for example by mechanical weed control, should be avoided since any kind of wounding increases the potential for invasion by *Phytophthora* spp. Soil fumigation is commonly used in bare root conifer nurseries to control *Phytophthora* spp., however it is not an economical practice in Christmas tree plantations or farms. Selective systemic fungicides are also used in nurseries, but fungicides are rarely effective in Christmas tree fields. In addition, from a Christmas tree grower's point of view, the practice of using fungicides in nurseries may not be beneficial since it does not cure the disease. It only suppresses the symptoms and may lead to extensive damage when asymptomatic plants are out in the field. The use of soil fumigants and other chemicals can be restricted. Please contact your national authorities to get information on the current issue on the national registration.

Quarantine recommendation

The European and Mediterranean Plant Protection Organization (EPPO) considers *P. ramorum* to be a dangerous organism. It is listed on the EPPO Alert List. For details see http://www.eppo.int/QUARANTINE/Alert_List/alert_list.htm.

In the European Union *P. ramorum* is a regulated organism according to the Commission Decision 2002/757/EU.

Literature used

- Adams, G.C., Bielenin, Jr. & A., 1988. First report of *Phytophthora cactorum* and *P. citricola*. Plant Disease 72:79.
- Benson, D.M., Grand, L.F., Suggs, E.G., 1976. Root rot of Fraser fir caused by *Phytophthora drechsleri*. Plant Dis. Rep. 60:238-240.
- Benson, D.M., Hinesley, L.E., Frampton, J., Parker, K.C., 1997. Evaluation of six *Abies* spp. To *Phytophthora* root rot caused by *Phytophthora cinnamomi*. APS Biological and Cultural Tests 13: 57.
- Chastagner, G.A., Riley, K.L., 2010. First report of *Phytophthora ramorum* infecting California Red Fir in California. Plant Disease 94: 1170.
- Chastagner, G.A., 2010. Susceptibility of true firs to *Phytophthora* root rot. A research update. Christmas Tree Outlook. P 23-27.
- Chastagner, G.A., Byther, R.S., 1997. *Phytophthora* root rot, stem canker, and shoot blight. Pages 28-30. In: G.A. Chastagner (Ed.). Christmas tree diseases, insects, & disorders in the Pacific Northwest: Identification and management. Washington State University, Cooperative Extension. 154 pp.
- Chastagner, G.A., Hamm, P.B., Riley, K.L., 1995. Symptoms and *Phytophthora* spp. associated with root rot and stem canker of noble fir Christmas trees in the Pacific Northwest. Plant Disease 79: 290-293.
- Cooley, S.J., Hamm, P.B., Hansen, E M., 1988. Susceptibility of Northwest conifers to *Phytophthora* root rot. Tree Planters' Notes 40:15-18.
- Frampton, J., Benson, D.M., 2012. Seedling resistance to *Phytophthora cinnamomi* in the genus *Abies*. Annals of Forest Science, Published On Line 8 May 2012. DOI 10.1007/s13595-012-0205-4
- Grand, L.F., Lapp, N.A., 1974. *Phytophthora cinnamomi* root rot of Fraser fir in North Carolina. Plant Dis. Rept. 58: 318-320.
- Hamm, P.B., Hansen, E.M., 1982. Pathogenicity of *Phytophthora* spp. to Northwest conifers. Eur. J. For. Pathology 12: 167-174.
- Hinesley, L.E., Parker, K.C., Benson, D.M., 2000. Evaluations of seedlings of Fraser, Momi, and Siberian fir for resistance to *Phytophthora cinnamomi*. HortScience 35:87-88.
- Hong, C.X., Marston, C.D., 2005. Crown rot of *Abies balsamea* var. *phanerolepis* caused by *Phytophthora cactorum* in Virginia. Plant Disease 89:433.
- Józsa, A., Z.Á. Nagy, A. Szigethy, G. Fischl, Bakonyi, J., 2011. First Report of *Phytophthora citrophthora* causing root and basal stem rot of woody ornamentals in Hungary. Plant Disease 95: 1193.
- McCain, A.H., Scharpf, R.F., 1986. *Phytophthora* shoot blight and canker disease of *Abies* spp. Plant Disease 70:1036-1037.
- Martin, F.N., Abad, Z.G., Balci, Y., Ivors, K., 2012. Identification and Detection of *Phytophthora*: Reviewing Our Progress, Identifying Our Needs. Plant Disease 96(8): 1080-1103.
- Quesada-Ocampo, L. M., Fulbright, D.W., Hausbeck, M.K., 2009. Susceptibility of Fraser fir to *Phytophthora capsici*. Plant Dis. 93:135-141.

Riley, K. L., G.A. Chastagner, Blomquist, C., 2011. First report of *Phytophthora ramorum* infecting grand fir in California. Plant Health Management, April. doi:10.1094/PHP-2011-0401-01-BR

Shew, H.D., Benson, D.M., 1981. Fraser fir root rot induced by *Phytophthora citricola*. Plant Dis. 65:688-689.

Talgø, V., M.L. Herrero, B. Toppe, S.S. Klemsdal, Stensvand, A., 2006. First report of root rot and stem canker caused by *Phytophthora cambivora* on noble fir (*Abies procera*) for bough production in Norway. Plant Disease 90(5):682.

Talgø, V., M.L. Herrero, B. Toppe, S.S. Klemsdal, Stensvand, A., 2007. *Phytophthora* root rot and stem canker found on Nordmann and subalpine fir in Norwegian Christmas tree plantations. Online. Plant Health Progress doi:10.1094/PHP-2007-0119-01-RS. 7 pp.

Links to further information

Phytophthora in the Forests:

<http://forestphytophthoras.org/>

P. ramorum:

www.suddenoakdeath.org

<http://rapra.csl.gov.uk/>

www.eppo.org

Phytophthora determination keys:

<http://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-08-11-0636>

Acknowledgement

This data sheet was prepared within the Working Group 1 of the European COST Action FP0801 (http://www.cost.eu/domains_actions/fps/Actions/FP0801).

Authors

Venche TALGØ¹, Gary CHASTAGNER²

¹ Norwegian Institute for Agricultural and Environmental Research, Plant Health and Plant Protection Division, Høgskoleveien 7, 1432 Ås, Norway
venche.talgo@bioforsk.no

² Washington State University, Research and Extension Center, 2606 West Pioneer Puyallup, WA 98371-4998, USA
chastag@wsu.edu

Disease symptoms of *Phytophthora* on *Abies* spec. (true firs)



Figure 1: Disease symptoms caused by root infection of *Phytophthora cambivora* on *Abies procera* (noble fir) in Norway (1)

Left: Discolored foliage

Right: Canker extending upwards from roots resulted in flagging and resin flow



Figure 2: Row of noble fir trees killed by *Phytophthora* root rot in a test plot at Washington State University

Abies procera (noble fir) are between two rows of *Abies nordmanniana* (Nordmann fir) (2)